MOLECULAR PHYSICS, 1991, VOL. 74, NO. 6, 1337–1343

Index of Authors

with the titles of papers

ACEVEDO, R., VASQUEZ, S. O., and FLINT, C. D.: Vibronic intensities in the electronic spectra of transition metal complex ions. Part IX. Experimental study of the relative vibronic intensities in the $^2E_g \rightarrow {}^4A_{2g}$ transition of the MnF_6^{2-} ion in Cs_2SiF_6, 843

ACEVEDO, R., VASQUEZ, S. O., and FLINT, C. D.: Vibronic intensities in the electronic spectra of transition metal complex ions. Part X. Second order Herzberg-Teller contributions to the vibronic intensity of the $^4A_{2g} \leftrightarrow {}^4T_{1g}$ transition in MnF_6^{2-}, 853

ADYA, A. K., and WORMALD, C. J.: First direct observation of orientationally correlated molecules in CO_2(1) by neutron diffraction, 735

ALEJANDRE, J., *see* Benavides, A. L.

ALLEN, J. M., ASHFORD, M. N. R., STICKLAND, R. J., and WESTERN, C. M.: The $\tilde{B}E''$ state of NH_3: the Jahn-Teller effect revealed by infrared-optical double resonance, 49

ALLEN, M. P., *see* Hess, S.

AMORETTI, G., *see* Caciuffo, R.

AMOS, M. D., and JACKSON, G.: BHS theory and computer simulations of linear heteronuclear triatomic hard-sphere molecules, 191

ANDRIESSEN, J., DORENBOS, P., and VAN EIJK, C. W. E.: Electronic structure and transition probabilities in pure and Ce^{3+} doped BaF_2, an explorative study, 535

AROUI, H., BROQUIER, M., CHEVALIER, M., PICARD, A., and BILLING, G. D.: Temperature effects on collision cross sections between some rovibrational states of ammonia gas perturbed by hydrogen and helium, 897

ASHFORD, M. N. R., *see* Allen, J. M.

ATTARD, P.: An improved Kirkwood superposition approximation for three atoms in rolling contact, 547

AURIEMMA, F., *see* Depner, M.

BACSKAY, G. B., *see* Wong, A. T.

BADIRKHAN, Z., PASTORE, G., and TOSI, M. P.: Charged hard spheres in a uniform neutralizing background: the role of thermodynamic self consistence, 1089

BAKER, J., *see* Barnes, M.

BARAM, A., and ROWLINSON, J. S.: Studies of the Gaussian model I. The one-component system, 707

BARANYAI, A., and EVANS, D. J.: Comments on thermodynamic integration methods for the determination of nonequilibrium entropy, 353

BARNES, M., BAKER, J., DYKE, J. M., FEHER, M., and MORRIS, A.: An investigation of the $SO(X^3\Sigma^-)$ molecule by multiphoton ionization spectroscopy, 689

BELLISSENT-FUNEL, M.-C., BUONTEMPO, U., PETRILLO, C., and RICCI, F. P.: Structure factor measurements of liquid Cl_2 at high temperature, 1209

BENAVIDES, A. L., ALEJANDRE, J., and DEL RIO, F.: Properties of the square-well fluid of variable width. IV. Molecular dynamics test of the van der Waals and long-range approximations, 321

BERMEL, W., *see* Ernst, M.

BHUIYAN, L. B., *see* Outhwaite, C. W.

BILLING, G. D., *see* Aroui, H.

BOCK, C. W., *see* Caminati, W.

BONDARENKO, G. V., and GORBATY, Y. E.: An infrared study of water vapour in the temperature range 573–723 K. Dimerization enthalpy and absorption intensities for monomer and dimer, 639

BOWDEN, G. J., MARTIN, J. P. D., and PRANDOLINI, M. J.: Non-square tensor algebra and application in the NMR of multiply connected spin systems: I, 985

BOWDEN, G. J., and PRANDOLINI, M. J.: Non-square tensor algebra applications in the NMR of multiply connected spin systems: II, 999
BRÄUCHLE, C., *see* Trentini, F. V.
BROQUIER, M., *see* Aroui, H.
BROWN, J. M., *see* Towle, J. P.
BRUNA, P. J., and GREIN, F.: Low-lying electronic states of Ge_2^+: a pseudopotential MRD-CI study, 1133
BRUSTOLON, M., *see* Caciuffo, R.
BUJÁN-NÚÑEZ, M. C., *see* López-Quintela, M. A.
BUONTEMPO, U., *see* Bellissent-Funel, M.-C.
BURGESS, A. N., *see* Hall, C. D.
BURIE, J-R., DAVIES, P. B., HANSFORD, G. M., MARTIN, N. A., GANG, J., and RUSSELL, D. K.: Diode laser infrared spectroscopy of jet-cooled hexacarbonyls of chromium, molybdenum, and tungsten, 919
BURNELL, E. E., *see* Ter Beek, L. C.

CACIUFFO, R., FRANCESCANGELI, O., GRECI, L., MELONE, S., GILLON, B., BRUSTOLON, M., MANIERO, A. L., AMORETTI, G., and SGARABOTTO, P.: Unpaired electron density in an indolinonic nitroxide radical as determined by polarized neutron diffraction, 905
CAMINATI, W., DAMIANI, D., CORBELLI, G., VELINO, B., and BOCK, C. W.: Microwave spectrum and *ab initio* calculations of ethylbenzene: potential energy surface of the ethyl group torsion, 885
CASTILLO, R., VILLAVERDE, A., and OROZCO, J.: Prediction of transport properties for Lennard-Jones fluids and their binary mixtures using the effective-diameter hard-sphere kinetic theory, 1315
CHALAM, M. K., *see* De Miguel, E.
CHEVALIER, R., *see* Aroui, H.
CHIERICO, A., *see* Mognaschi, E. R.
CHILD, M. S., *see* Gilbert, R. D.
CORBELLI, G., *see* Caminati, W.

DAHM, J., MACHO, V., MARESCH, G. G., and SPIESS, H. W.: Photopolymerization study of photoresists by ESR spectroscopy and ESR imaging, 591
DAMIANI, D., *see* Caminati, W.
DAVIES, P. B., *see* Burie, J-R.
DE MIGUEL, E., RULL, L. F., CHALAM, M. K., and GUBBINS, K. E.: Liquid crystal phase diagram of the Gay-Berne fluid, 405
DEL RIO, F., *see* Benavides, A. L.
DEITERS, U. K.: The equation of state of soft repulsive spherical molecules, 153
DEPNER, M., SCHÜRMANN, B. L., and AURIEMMA, F.: Investigation of a poly(oxyethylene) chain by a molecular dynamics simulation in an aqueous solution and by Langevin dynamics simulations, 715
DICKINSON, E., and PELAN, E. G.: Molecular dynamics model of adsorbed layer at a plane fluid interface, 1115
DOLG, M., KÜCHLE, W., STOLL, H., PREUSS, H., and SCHWERDTFEGER, P.: *Ab initio* pseudopotentials for Hg to Rn. II. Molecular calculations on the hydrides of Hg to At and the fluorides of Rn, 1265
DOLG, M., *see* Küchle, W.
DOMINGUEZ-ADAME, F., MÉNDEZ, B., MACIÁ, E., and GONZALEZ, M. A.: Non-local separable potential approach to multicentre interactions, 1065
DORENBOS, P., *see* Andriessen, J.
DUNCAN, J. L., *see* Law, M. M.
DYKE, J. M., *see* Barnes, M.
DYKE, J. M., *see* Lee, E. P. F.

ENO, L.: End-point sensitivity in quantum dynamic calculations, 923
ERNST, M., GRIESINGER, C., ERNST, R. R., and BERMEL, W.: Optimized heteronuclear cross polarization in liquids, 219

ERNST, R. R., *see* Ernst, M.
ESTEBAN, A. L., and GALACHE, M. P.: The r_z structure of cyclobutane from its dipolar couplings, 283
EVANS, D. J., *see* Baranyai, A.
EVANS, G. T., and MADDEN, P. A.: Vibrational energy transfer and pair dynamics in liquids: a kinetic theory study, 1171
EVANS, G. T., and SMITH, E. B.: Second virial coefficients for nonspherical molecules: a 'convex peg in a round hole' potential, 79
EVANS, G. T.: Transport coefficients of nonspherical hard bodies: a kinetic theory approach, 775
EVANS, G. T., *see* Tjipto-Margo, B.

FEHER, M., *see* Barnes, M.
FILABOZZI, A., and POSTORINO, P.: New determination of the He–Xe gas–gas demixing surface in the high Xe concentration region, 273
FLINT, C. D., *see* Acevedo, R.
FOWLER, P. W., and STEINER, E.: Paramagnetic closed-shell molecules: the isoelectronic series CH^+, BH and BeH^-, 1147
FRANCESCANGELI, O., *see* Caciuffo, R.
FRENKEL, D., *see* Hess, S.
FRENKEL, D., *see* Mooij, G. C. A. M.
FRENKEL, D., *see* Smit, B.
FRONEMAN, S., *see* Reynhardt, E. C.
FUSHIKI, M.: A triplet correlation function for Lennard-Jones fluids, 307

GALACHE, M. P., *see* Esteban, A. L.
GALLEGO, L. J., REY, C., and GRIMSON, M. J.: A Monte Carlo simulation study of the disjoining pressure in thin fluid films sterically stabilized by terminally attached chains, 383
GANG, J., *see* Burie, J-R.
GIANTURCO, F. A., SERNA, S., and SANNA, N.: Dynamical decoupling in the quantum calculations of transport coefficients. I—coupled state results for He–N_2 gaseous mixture, 1071
GILBERT, R. D., CHILD, M. S., and JOHNS, J. W. C.: Rotational analysis and assignment of the $3d \leftarrow 1b_1$ Rydberg complex in H_2O and D_2O, 473
GILLON, B., *see* Caciuffo, R.
GLAZ, W.: On 'cross' spectral moments of the collisional polarizabilities of rigid rotors, 1019
GONZALEZ, D. J., GONZALEZ, L. E., and SILBERT, M.: Structure and thermodynamics of hard *D*-dimensional spheres: overlap volume function approach, 613
GONZALEZ, L. E., *see* Gonzalez, D. J.
GONZALEZ, M. A., *see* Dominguez-Adame, F.
GORBATY, Y. E., *see* Bondarenko, G. V.
GORDALLA, B. C., and ZEIDLER, M. D.: NMR proton relaxation and chemical exchange in the system $H_2^{16}O/H_2^{17}O-|^2H_6|$ dimethylsulphoxide, 975
GRECI, L., *see* Caciuffo, R.
GREEN, N. J. B., and PIMBLOTT, S. M.: Scavenger kinetics and the Laplace transform relationship. 1. Homogeneous systems, 795
GREEN, N. J. B., and PIMBLOTT, S. M.: Scavenger kinetics and the Laplace transform relationship. 2. Nonhomogeneous kinetics, 811
GREIN, F., *see* Bruna, P. J.
GREENSPAN, D., On electron attraction, 1015
GRIESINGER, C., *see* Ernst, M.
GRIMSON, M. J., and SILBERT, M.: A self-consistent theory of the effective interactions in charge-stabilized colloidal dispersions, 397
GRIMSON, M. J.: Structure and elasticity of model gel networks, 1097
GRIMSON, M. J., *see* Gallego, L. J.
GUBBINS, K. E., *see* De Miguel, E.

HALL, C. D., JOHNSON, K. A., BURGESS, A. N., WINTERTON, N., and HOWELLS, W. S.: The structure of fluid fluoroform, chlorodifluoromethane, and dichlorodifluoromethane by neutron diffraction, 27
HAMLEY, I. W., LUCKHURST, G. R., and SEDDON, J. M.: The influence of molecular biaxiality on intramolecular X-ray and neutron scattering from uniaxial nematic liquid crystals, 1221
HANSFORD, G. M., *see* Burie, J-R.
HANSON, E. E., *see* Wong, C. C. K.
HARADA, M., *see* Kinoshita, M.
HENDERSON, J. R.: A derivation of external fields that create step function density profiles at wall–fluid interfaces, 1125
HESS, S., FRENKEL, D., and ALLEN, M. P.: On the anisotropy of diffusion in nematic liquid crystals: test of a modified affine transformation model via molecular dynamics, 765
HOFFMANN, H., *see* Rehage, H.
HOHM, U.: A new method for the determination of the dispersion of the second recractivity virial coefficient $b_R(\lambda)$ and its application to Ar, Kr, and CO_2, 1233
HOWELLS, W. S., *see* Hall, C. D.
HUSH, N. S., *see* Wong, A. T.

IMRIE, D. A., and RAAB, R. E.: A new molecular theory of field gradient induced birefringence used for measuring electric quadrupole moments, 833

Jackson, G., *see* Amos, M. D.
JACOBS, P. W. M., *see* Rycerz, Z. A.
JOHNS, J. W. C., *see* Gilbert, R. D.
JOHNSON, K. A., *see* Hall, C. D.

KENYON, A. J., MCCAFFERY, A. J., QUINTELLA, C. M., and WINKEL, J. F.: A study of molecular dynamics within liquid flows using fluorescence depolarization, 871
KINOSHITA, M., and HARADA, M.: Numerical solution of the HNC equation for fluids of non-spherical particles. An efficient method with application to dipolar hard spheres, 443
KINOSHITA, M., HARADA, M., and SHIOI, A.: Singularity of the HNC equation occuring at a very high density for hard spheres, 927
KJELLANDER, R., SARMAN, S.: A study of anisotropic pair distribution theories for Lennard-Jones fluids in narrow slits. Part II. Pair correlations and solvation forces, 665
KOLAFA, J.: Nebular boundary conditions in Monte Carlo simulations, 143
KONDO, A. E., and MEATH, W. J.: On the dynamics of pulsed and continuous wave laser-molecule interactions and the effects of permanent dipoles, 113
KOWALEWSKI, J., *see* Ribbing, C.
KREMP, D., *see* Sonnenburg, J.
KÜCHLE, W., DOLG, M., STOLL, H., and PREUSS, H.: *Ab initio* pseudopotentials for Hg through Rn. I. Parameter sets and atomic calculations, 1245
KÜCHLE, W., *see* Dolg, M.

LAAKSONEN, A., and STILBS, P.: Molecular dynamics and NMR study of methane–water systems, 747
LABÍK, S., MALIJEVSKÝ, A., POSPÍŠIL, R., and SMITH, W. R.: An accurate integral equation for molecular fluids. IV. Hard prolate spherocylinders, 261
LABÍK, S., *see* Pospíšil, R.
LABORANTI, L. M., *see* Mognaschi, E. R.
LAGO, S., *see* Padilla, P.
LAW, M. M., and DUNCAN, J. L.: C–H and C–D interbond coupling and Fermi resonance interactions in CHD_2Cl, 861
LEE, E. P. F., and DYKE, J. M.: An *ab initio* molecular orbital study of the anions $O_2^- \cdot H_2O$ and $O_2^- \cdot CO_2$, 333
LEVESQUE, D., *see* Zarragoicoechea, G. J.
LÓPEZ-QUINTELA, M. A., TOJO, C., and BUJÁN-NÚÑEZ, M. C.: 'Slow-down' of diffusion coefficient in finite Brownian motion, 785
LUCKHURST, G. R., *see* Hamley, I. W.

MACHO, V., *see* Dahm, J.
MACIÁ, E., *see* Dominguez-Adame, F.
MADDEN, P. A., *see* Evans, G. T.
MALIJEVSKÝ, A., *see* Labík, S.
MALIJEVSKÝ, A., *see* Pospíšil, R.
MANIERO, A. L., *see* Caciuffo, R.
MARESCH, G. G., *see* Dahm, J.
MAROULIS, G.: On the dipole polarizability of HCl, 131
MARTIN, J. P. D., *see* Bowden, G. J.
MARTIN, N. A., *see* Burie, J-R.
MASTERS, A. J., *see* Treloar, R. L.
MCCAFFERY, A. J., *see* Kenyon, A. J.
MCCOURT, F. R. W., *see* Wong, C. C. K.
MEATH, W. J., *see* Kondo, A. E.
MELONE, S., *see* Caciuffo, R.
MÉNDEZ, B., *see* Dominguez-Adame, F.
MOGNASCHI, E. R., LABORANTI, L. M., and CHIERICO, A.: On the correlation factor of pure polar fluids whose molecules dimerize to nonpolar dimers, 211
MOOIJ, G. C. A. M., and FRENKEL, D.: Novel scheme to compute chemical potentials of chain molecules on a lattice, 41
MORRIS, A., *see* Barnes, M.

ODELIUS, M., *see* Ribbing, C.
ORLIKOWSKI, T., *see* Wolniewicz, L.
OROZCO, J., *see* Castillo, R.
OUTHWAITE, C. W., and BHUIYAN, L. B.: A modified Poisson-Boltzmann analysis of the electric double layer around an isolated spherical macroion, 367

PADILLA, P., LAGO, S., and VEGA, C.: A bridge between Fischer and Boublik thermodynamic perturbation theories: calculating thermodynamic properties of pure substances, 161
PASTORE, G., *see* Badirkhan, Z.
PELAN, E. G., *see* Dickinson, E.
PERIĆ, M., *see* Reuter, W.
PETRI, J., *see* Castillo, R.
PETRILLO, C., *see* Bellissent-Funel, M.-C.
PEYERIMHOFF, S. D., *see* Reuter, W.
PICARD, A., *see* Aroui, H.
PIMBLOTT, S. M., *see* Green, N. J. B.
POSPÍŠIL, R., MALIJEVSKÝ, A., LABÍK, S., and SMITH, W. R.: An accurate integral equation for molecular fluids. III. Hard linear homonuclear triatomics, 253
POSPÍŠIL, R., *see* Labík, S.
POSTORINO, P., *see* Filabozzi, A.
PRANDOLINI, M. J., *see* Bowden, G. J.
PREUSS, H., *see* Dolg, M.
PREUSS, H., *see* Küchle, W.

QUINTELLA, C. M., *see* Kenyon, A. J.

RAAB, R. E., *see* Imrie, D. A.
REHAGE, H., and HOFFMANN, H.: Viscoelastic surfactant solutions: model systems for rheological research, 933
REUTER, W., PERIĆ, M., and PEYERIMHOFF, S. D.: *Ab initio* study of the vibronic structure for the X^2B_1 and A^2A_1 electronic states of H_2O^+, 569
RIBBING, C., ODELIUS, M., and KOWALEWSKI, J.: Simple nonempirical calculations of the zero field splitting in transition metal systems. II. NiF_6^{4-} and Ni(II) complexes with mixed ligands, 1299
RICCI, F. P., *see* Bellissent-Funel, M.-C.
ROWLINSON, J. S., *see* Baram, A.
REY, C., *see* Gallego, L. J.

REYNHARDT, E. C., and FRONEMAN, S.: Phase transitions and molecular motions in crystalline *ortho-* and *meta-* carborane, 61
RUDOWICZ, C.: Correlations between orthorhombic crystal field parameters for rare-earth (f^n) and transition-metal (d^n) ions in crystals: $REBa_2Cu_3O_{7-x}$, $RE_2F_{14}B$, RE-garnets, RE: LaF_3 and MnF_2, 1159
RULL, L. F., *see* De Miguel, E.
RUSSELL, D. K., *see* Burie, J-R.
RYCERZ, Z. A., and JACOBS, P. W. M.: Effective pair potentials in liquid metals. A comparative study for liquid lead, 425

SÄNDIG, R., *see* Sonnenburg, J.
SANNA, N., *see* Gianturco, F. A.
SARMAN, S., *see* Kjellander, R.
SCHÜRMANN, B. L., *see* Depner, M.
SCHWERDTFEGER, P., *see* Dolg, M.
SEDDON, J. M., *see* Hamley, I. W.
SERNA, S., *see* Gianturco, F. A.
SESE, L. M.: Path-integral and effective potential Monte Carlo simulations of liquid nitrogen, hard-sphere and Lennard-Jones potentials, 177
SGARABOTTO, P., *see* Caciuffo, R.
SHINOHARA, T., *see* Sogami, I. S.
SHIOI, A., *see* Kinoshita, M.
SILBERT, M., *see* Gonzalez, D. J.
SILBERT, M., *see* Grimson, M. J.
SIMON, U., *see* Zimmermann, W.
SMALLEY, M. V., *see* Sogami, I. S.
SMIT, B., and FRENKEL, D.: Vapour-liquid equilibria of the hard core Yukawa fluid, 35
SMITH, E. B., *see* Evans, G. T.
SMITH, W. R., *see* Labík, S.
SMITH, W. R., *see* Pospíšil, R.
SOGAMI, I. S., SHINOHARA, T., and SMALLEY, M. V.: Effective interaction of highly charged plates in an electrolyte, 599
SONNENBURG, J., KREMP, D., and SÄNDIG, R.: Diffusion processes in Brownian particle systems with short-range interaction, 649
SPIESS, H. W., *see* Dahm, J.
STEINER, E., *see* Fowler, P. W.
STICKLAND, R. J., *see* Allen, J. M.
STILBS, P., *see* Laaksonen, A.
STOLL, H., *see* Dolg, M.
STOLL, H., *see* Küchle, W.

TELEMAN, O., *see* Wallqvist, A.
TER BEEK, L. C., ZIMMERMAN, D. S., and BURNELL, E. E.: The conformation of 2,2′-dithiophene in nematic solvents determined by ^{1}H–NMR, 1027
TJIPTO-MARGO, B., and EVANS, G. T.: A van der Waals theory of nematic liquid crystals: a 'convex peg in a round hole' potential, 85
TOJO, C., *see* López-Quintela, M. A.
TOSI, M. P., *see* Badirkhan, Z.
TOWLE, J. P., and BROWN, J. M.: The infrared spectrum of the TeD radical by laser magnetic resonance, 465
TRELOAR, R. L., and MASTERS, A. J.: Short-time diffusion of anisotropic particles in suspension —a perturbative approach, 1195
TRENTINI, F. W., VOITLÄNDER, J., and BRÄUCHLE, C.: Resonance theoretical description of magnetic zero field splitting energies of benzenoid hydrocarbons. Structural dependence of the parameter *D*, 553

URBAN, W., *see* Zimmermann, W.

VAN EIJK, C. W. E., *see* Andriessen, J.
VASQUEZ, S. O., *see* Acevedo, R.
VEGA, C., *see* Padilla, P.
VELINO, B., *see* Caminati, W.
VILLAVERDE, A., *see* Castillo, R.
VOITLÄNDER, J., *see* Trentini, F. V.

WALES, D. J.: Theoretical study of some small van der Waals complexes containing inert gas atoms, 1
WALLQVIST, A., and TELEMAN, O.: Properties of flexible water models, 515
WEBER, J., and WOKAUN, A.: Two-dimensional NMR studies of the conformational B → Z transition in the oligodeoxynucleotides $d(CG)_8$ and $d(m^5CG)_8$, 293
WEIS, J. J., *see* Zarragoicoechea, G. J.
WESTERN, C. M., *see* Allen, J. M.
WINKEL, J. F., *see* Kenyon, A. J.
WINTERTON, N., *see* Hall, C. D.
WOKAUN, A., *see* Weber, J.
WOLNIEWICZ, L., and ORLIKOWSKI, T.: The $1s\sigma_g$ and $2p\sigma_u$ states of the H_2^+, D_2^+ and HD^+ ions, 103
WONG, A. T., BACSKAY, G. B., and HUSH, N. S.: *Ab initio* polarizability derivatives for methane: an application to Raman intensities of overtone and combination bands, 1037
WONG, C. C. K., HANSON, E. E., and MCCOURT, F. R. W.: Classical trajectory calculation of transport and relaxation properties for CO_2–He mixtures, 497
WORMALD, C. J., *see* Adya, A. K.

ZARRAGOICOECHEA, G. J., LEVESQUE, D., and WEIS, J. J.: Monte Carlo study of dipolar ellipsoids. I. The isotropic phase, 629
ZEIDLER, M. D., *see* Gordalla, B. C.
ZIMMERMAN, D. S., *see* Ter Beek, L. C.
ZIMMERMANN, W., SIMON, U., PETRI, M., and URBAN, W.: The observation of fine-structure transitions in the SnH radical by diode laser spectroscopy, 1287